



Lignin Valorization by Heterogeneous Catalytic Oxidation

Melián Rodríguez, Mayra; Shunmugavel, Saravanamurugan; Kegnæs, Søren; Riisager, Anders

Publication date:
2015

[Link back to DTU Orbit](#)

Citation (APA):

Melián Rodríguez, M., Shunmugavel, S., Kegnæs, S., & Riisager, A. (2015). *Lignin Valorization by Heterogeneous Catalytic Oxidation*. Abstract from 4th International Congress in Green Process Engineering, Sevilla, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LIGNIN VALORIZATION BY HETEROGENEOUS CATALYTIC OXIDATION

M. MELIÁN-RODRÍGUEZ (1), S. SHUNMUGAVEL (1), S. KEGNÆS (1), A. RIISAGER*(1)

♦ Centre for Catalysis and Sustainable Chemistry
Technical University of Denmark, Department of Chemistry
Kemitorvet, building 207
DK-2800 Kgs. Lyngby, Denmark
e-mail : mayro@kemi.dtu.dk

Key-words. lignin, green chemistry, biomass, heterogeneous catalysis, oxidation

Many different technologies to convert biomass into high-value products and fuels in “bio-refineries” have been proposed in the literature to meet present and approaching future challenges such as, global warming, high oil prices and food shortage. In this context, special attention is giving to the conversion of lignocellulosic biomass, which does not compete with food resources and is widely available as a low cost feedstock. Lignocellulose biomass is composed of three main fractions; cellulose (40-50 wt.%), hemicellulose (25-35 wt.%) and lignin (20-30 wt.%). Lignin represents 40% of the energy content of lignocellulosic biomass, and it is the obvious candidate to serve as an available feedstock for the production of basic aromatic chemicals [1].

In green plants the molecular weight and amount of lignin differ, but it is mainly composed of three different monolignol monomers: *p*-coumaryl, coniferyl and sinapyl alcohol [2]. These monomers are connected with various linkages with the most common one being the β -O-4 linkage (Figure 1) [3].

The lignin structure is complex so different model compounds are often used to study lignin valorization. These model compounds contain the linkages present in lignin, simplifying catalytic analysis and present analytical challenges related to the study of the complicated lignin polymer and the plethora of products that could be obtained [2].




Figure 1. Linkages present in lignin.

For valorization of lignin (and model compounds) heterogeneous oxidative catalysis plays an important role due to the industrial viability of such systems [4]. Lange et al. have reported that the catalytic oxidation products of lignin and lignin model compounds can range from aromatic aldehydes to carboxylic acid originating from oxidation of side chains depending on the reaction conditions [5].

In this work, we have examined the oxidative conversion of several lignin model compounds using numerous heterogeneous catalysts and oxygen as oxidant.

REFERENCES

- [1] P.T. Patil, U. Armbruster et al., Energy Fuels, 25 (2011) 4713.
- [2] J. Zakzeski, P.C.A Bruijninx et al., Chem Rev, 110 (2010) 3552
- [3] P. T. Patil et al., Energy Fuels, 25 (2011) 4713
- [4] Q. Xiang, Y.Y. Lee, Appl. Biochem. Biotechnol., 153 (2000) 84
- [5] H. Lange et al., Eur. Polymer J., 49 (2013) 1151